

demographic factors such a bottleneck associated with human migration out of Africa 50,000 to 100,000 years ago.

Previous studies have shown that *Microcephalin* is a specific regulator of brain size (13, 14) and that this gene has evolved under strong positive selection in the primate lineage leading to *Homo sapiens* (7, 8). Here, we present compelling evidence that *Microcephalin* has continued its trend of adaptive evolution beyond the emergence of anatomically modern humans. The specific function of *Microcephalin* in brain development makes it likely that selection has operated on the brain. Yet, it remains formally possible that an unrecognized function of *Microcephalin* outside of the brain is actually the substrate of selection. If selection indeed acted on a brain-related phenotype, there could be several possibilities, including brain size, cognition, personality, motor control, or susceptibility to neurological and/or psychiatric diseases. We hypothesize that D and non-D haplotypes have different effects on the proliferation of neural progenitor cells, which in turn leads to different phenotypic outcomes of the brain visible to selection.

References and Notes

1. J. N. Spuhler, *The Evolution of Man's Capacity for Culture* (Wayne State Univ. Press, Detroit, MI, 1959).
2. J. H. Jerison, *Evolution of the Brain and Intelligence* (Academic Press, New York, 1973).
3. W. Enard et al., *Nature* **418**, 869 (2002).

4. J. Zhang, *Genetics* **165**, 2063 (2003).
5. P. D. Evans et al., *Hum. Mol. Genet.* **13**, 489 (2004).
6. N. Kouprina et al., *PLoS Biol.* **2**, E126 (2004).
7. Y. Q. Wang, B. Su, *Hum. Mol. Genet.* **13**, 1131 (2004).
8. P. D. Evans, J. R. Anderson, E. J. Vallender, S. S. Choi, B. T. Lahn, *Hum. Mol. Genet.* **13**, 1139 (2004).
9. R. J. Ferland et al., *Nat. Genet.* **36**, 1008 (2004).
10. H. H. Stedman et al., *Nature* **428**, 415 (2004).
11. F. Burki, H. Kaessmann, *Nat. Genet.* **36**, 1061 (2004).
12. S. Dorus et al., *Cell* **119**, 1027 (2004).
13. A. P. Jackson et al., *Am. J. Hum. Genet.* **63**, 541 (1998).
14. A. P. Jackson et al., *Am. J. Hum. Genet.* **71**, 136 (2002).
15. W. B. Dobyns, *Am. J. Hum. Genet.* **112**, 315 (2002).
16. G. H. Mochida, C. A. Walsh, *Curr. Opin. Neurol.* **14**, 151 (2001).
17. C. G. Woods, J. Bond, W. Enard, *Am. J. Hum. Genet.* **76**, 717 (2005).
18. E. Roberts et al., *Eur. J. Hum. Genet.* **7**, 815 (1999).
19. L. Moynihan et al., *Am. J. Hum. Genet.* **66**, 724 (2000).
20. C. R. Jamieson, C. Govaerts, M. J. Abramowicz, *Am. J. Hum. Genet.* **65**, 1465 (1999).
21. J. Bond et al., *Nat. Genet.* **32**, 316 (2002).
22. G. F. Leal et al., *J. Med. Genet.* **40**, 540 (2003).
23. J. Bond et al., *Nat. Genet.* **37**, 353 (2005).
24. T. Huyton, P. A. Bates, X. Zhang, M. J. Sternberg, P. S. Freemont, *Mutat. Res.* **460**, 319 (2000).
25. S. Y. Lin, S. J. Eledge, *Cell* **113**, 881 (2003).
26. X. Xu, J. Lee, D. F. Stern, *J. Biol. Chem.* **279**, 34091 (2004).
27. M. Trimborn et al., *Am. J. Hum. Genet.* **75**, 261 (2004).
28. S. L. Gilbert, W. B. Dobyns, B. T. Lahn, *Nat. Rev. Genet.* **6**, 581 (2005).
29. M. Stephens, N. J. Smith, P. Donnelly, *Am. J. Hum. Genet.* **68**, 978 (2001).
30. M. Stephens, P. Donnelly, *Am. J. Hum. Genet.* **73**, 1162 (2003).
31. S. A. Tishkoff et al., *Science* **293**, 455 (2001).
32. P. C. Sabeti et al., *Nature* **419**, 832 (2002).
33. E. Wang et al., *Am. J. Hum. Genet.* **74**, 931 (2004).
34. T. Bersaglieri et al., *Am. J. Hum. Genet.* **74**, 1111 (2004).
35. E. E. Thompson et al., *Am. J. Hum. Genet.* **75**, 1059 (2004).
36. H. Stefansson et al., *Nat. Genet.* **37**, 129 (2005).

37. R. R. Hudson, *Oxf. Surv. Evol. Biol.* **7**, 1 (1990).
38. R. R. Hudson, *Bioinformatics* **18**, 337 (2002).
39. A. Kong et al., *Nat. Genet.* **31**, 241 (2002).
40. E. Zietkiewicz et al., *J. Mol. Evol.* **47**, 146 (1998).
41. H. Harpending, A. Rogers, *Annu. Rev. Genomics Hum. Genet.* **1**, 361 (2000).
42. H. Tang, D. O. Siegmund, P. Shen, P. J. Oefner, M. W. Feldman, *Genetics* **161**, 447 (2002).
43. R. Thomson, J. K. Pritchard, P. Shen, P. J. Oefner, M. W. Feldman, *Proc. Natl. Acad. Sci. U.S.A.* **97**, 7360 (2000).
44. I. McDougall, F. H. Brown, J. G. Fleagle, *Nature* **433**, 733 (2005).
45. R. G. Klein, *The Human Career: Human Biological and Cultural Origins* (Univ. of Chicago Press, Chicago, 1999).
46. M. Bamshad, S. P. Wooding, *Nat. Rev. Genet.* **4**, 99 (2003).
47. F. Tajima, *Genetics* **123**, 585 (1989).
48. S. Wright, *Evolution and the Genetics of Populations* (Univ. of Chicago Press, Chicago, 1978).
49. J. M. Akey, G. Zhang, K. Zhang, L. Jin, M. D. Shriver, *Genome Res.* **12**, 1805 (2002).
50. We thank the Coriell Institute for Medical Research, the Centre d'Etude du Polymorphisme Humain (CEPH), and A. Froment for human DNA samples. We thank H. M. Cann, S. Dorus, E. E. Eichler, N. M. Pearson, A. Di Rienzo, M. Kreitman, and J. K. Pritchard for technical support and/or helpful discussions. Supported in part by the Searle Scholarship and the Burroughs Wellcome Career Award (to B.T.L.), and David and Lucile Packard Career Award, the Burroughs Wellcome Career Award, and NSF grant BCS-0196183 (to S.A.T.).

Supporting Online Material

www.sciencemag.org/cgi/content/full/309/5741/1717/DC1  
 Materials and Methods  
 Tables S1 and S2  
 References and Notes

18 April 2005; accepted 14 June 2005  
 10.1126/science.1113722

# Ongoing Adaptive Evolution of *ASPM*, a Brain Size Determinant in *Homo sapiens*

Nitzan Mekel-Bobrov,<sup>1,2</sup> Sandra L. Gilbert,<sup>1</sup> Patrick D. Evans,<sup>1,2</sup> Eric J. Vallender,<sup>1,2</sup> Jeffrey R. Anderson,<sup>1</sup> Richard R. Hudson,<sup>3</sup> Sarah A. Tishkoff,<sup>4</sup> Bruce T. Lahn<sup>1\*</sup>

The gene *ASPM* (*abnormal spindle-like microcephaly associated*) is a specific regulator of brain size, and its evolution in the lineage leading to *Homo sapiens* was driven by strong positive selection. Here, we show that one genetic variant of *ASPM* in humans arose merely about 5800 years ago and has since swept to high frequency under strong positive selection. These findings, especially the remarkably young age of the positively selected variant, suggest that the human brain is still undergoing rapid adaptive evolution.

Homozygous null mutations of *ASPM* cause primary microcephaly, a condition characterized by severely reduced brain size with otherwise normal neuroarchitecture (1). Studies


have suggested that *ASPM* may regulate neural stem cell proliferation and/or differentiation during brain development, possibly by mediating spindle assembly during cell division (1, 2). Phylogenetic analysis of *ASPM* has revealed strong positive selection in the primate lineage leading to *Homo sapiens* (3–5), especially in the past 6 million years of hominid evolution in which *ASPM* acquired about one advantageous amino acid change every 350,000 years (4). These data argue that *ASPM*

may have contributed to human brain evolution (3–6). Here, we investigate whether positive selection has continued to operate on *ASPM* since the emergence of anatomically modern humans.

Human *ASPM* has 28 exons with a 10,434–base pair open reading frame (1) (fig. S1). We resequenced the entire 62.1-kb genomic region of *ASPM* in samples from 90 ethnically diverse individuals obtained through the Coriell Institute and from a common chimpanzee (7). This revealed 166 polymorphic sites (table S1). Using established methodology (7), we identified 106 haplotypes. One haplotype, numbered 63, had an unusually high frequency of 21%, whereas the other haplotypes ranged from 0.56% to 3.3% (fig. S2). Moreover, this haplotype differed consistently from the others at multiple polymorphic sites (save for a few rare haplotypes that are minor mutational or recombinational variants of haplotype 63, as discussed later) (table S2). Two of these polymorphic sites are nonsynonymous, both in exon 18, and are denoted A44871G and C45126A (numbers indicate genomic positions from the start codon, and letters at the beginning and end indicate ancestral and derived alleles, respectively). These two sites reside in a region of the open reading frame that was shown previously to have experienced par-

<sup>1</sup>Howard Hughes Medical Institute, Department of Human Genetics, <sup>2</sup>Committee on Genetics, <sup>3</sup>Department of Ecology and Evolution, University of Chicago, Chicago, IL 60637, USA. <sup>4</sup>Department of Biology, University of Maryland, College Park, MD 20742, USA.

\*To whom correspondence should be addressed. E-mail: blahn@bsd.uchicago.edu



**Fig. 1.** Worldwide frequencies of *ASPM* haplogroup D chromosomes (defined as having the derived G allele at the A44871G diagnostic polymorphism), based on a panel of 1186 individuals.

ticularly strong positive selection in the lineage leading to humans (4) (fig. S1).

The unusually high frequency of haplotype 63 is strongly suggestive of positive selection (8). We tested the statistical significance of the possibility of positive selection using coalescent modeling (7). The frequency of haplotype 63 is notably higher in Europeans and Middle Easterners (including Iberians, Basques, Russians, North Africans, Middle Easterners, and South Asians), as compared with other populations (table S1). We therefore focused on this group to take advantage of its relatively simple and homogeneous demographic structure (9). Because 7 of the 50 Europeans and Middle Easterners were homozygous for haplotype 63, we tested the probability of obtaining 7 or more homozygotes (among 50) for a single haplotype across a 62.1-kb region containing 122 segregating sites (the number of polymorphic sites found in Europeans and Middle Easterners). The recombination rate and the gene conversion rate of the locus used in the test were obtained from our polymorphism data (7). For demographic history, we

assumed a severe bottleneck followed by exponential growth (7) that is likely to be much more stringent than the bottleneck associated with the colonization of Eurasia (10). These parameters produced a highly significant departure from the neutral expectation ( $P < 0.00001$ ). The simulation was repeated with a wide range of demographic histories (7), all of which produced highly significant results. We repeated the above tests on the entire Coriell panel, which also demonstrated strong significance. Finally, we repeated these tests using the inferred frequency of haplotype 63 (instead of the number of individuals who are homozygous for this haplotype) and again obtained significant statistical values. These data indicate that haplotype 63 has spread to high frequency under positive selection. The two nonsynonymous polymorphisms previously mentioned are either the target of selection or closely linked to the target site.

We define haplogroup D (where D stands for “derived”) as the class of haplotypes with the derived G allele at the A44871G nonsynonymous polymorphic site. The haplotypes

with the A allele are defined as non-D. This classification is meant to capture the notable structure in the 106 haplotypes. Haplogroup D comprises two subgroups. One contains the predominant haplotype 63 and its closely related mutational variants [including haplotypes 64, 66, and 71 (table S2)]. The other contains recombinant haplotypes between haplotype 63 (or its close mutational variants) and non-D haplotypes (including haplotypes 62, 65, and 105). The frequency of haplogroup D chromosomes is 28% in the entire Coriell panel and 44% in Europeans and Middle Easterners. Another prominent feature separating D and non-D haplotypes is the fact that the two classes have fixed differences relative to each other at multiple sites, where D haplotypes have the derived alleles (except for the rare recombinants between D and non-D chromosomes). This unusual haplotype structure is consistent with the following evolutionary history: (i) a rapid increase of haplotype 63 from a single ancestral copy to high frequency, and (ii) the introduction of minor variants of haplotype 63 by mutation and by recombination between D

and non-D chromosomes (these variants make up the other members of haplogroup D).

Another signature of positive selection is extended linkage disequilibrium (LD) (8), which is evident in the 62.1-kb region. Indeed, the 50 haplogroup D chromosomes show nearly complete LD across the region, with only three cases of LD breakdown [haplotypes 62, 65, and 105, present in one, two, and one copy, respectively (table S2)]. In contrast, the non-D chromosomes do not show any unusual LD across the region. We investigated the decay of LD beyond the 62.1-kb region by sequencing the Coriell panel for two flanking intergenic segments of roughly 5 kb each, positioned 25 kb upstream and downstream of *ASPM*. There is notable LD breakdown in these two segments, confirming that the site of selection is most likely within *ASPM*.

Following established methodology (7), we estimated the coalescence age (i.e., time to the most recent common ancestor) of haplogroup D at 5800 years, with a 95% confidence interval between 500 and 14,100 years. In comparison, the coalescence age of all the chromosomes (both D and non-D) is ~800,000 years. Thus, the age of haplogroup D substantially postdates the emergence of anatomically modern humans, estimated at ~200,000 years ago (11). A rough calculation showed that the fitness advantage of haplogroup D is in the range of a few percent over non-D haplotypes.

By genotyping the A44871G diagnostic polymorphism described earlier, we obtained the global frequency distribution of haplogroup D chromosomes from a separate panel of 1186 individuals (7). Consistent with the

Coriell panel, we observed much higher frequency of haplogroup D chromosomes in Europeans and Middle Easterners than in other populations (Fig. 1). The corresponding estimate of  $F_{ST}$ , a statistic of genetic differentiation, is 0.29 between Europeans/Middle Easterners and other populations and 0.31 between Europeans/Middle Easterners and sub-Saharan Africans. These values indicate considerable genetic differentiation at this locus (12). Several scenarios may account for such notable differentiation. One is that haplogroup D first arose somewhere in Eurasia and is still in the process of spreading to other regions. The other is that it arose in sub-Saharan Africa, but reached higher frequency outside of Africa partly because of the bottleneck during human migration out of Africa. Finally, it is possible that differential selective pressure in different geographic regions is partly responsible.

Collectively, our data offer strong evidence that haplogroup D emerged very recently and subsequently rose to high frequency under strong positive selection. The recent selective history of *ASPM* in humans thus continues the trend of positive selection that has operated at this locus for millions of years in the hominid lineage (3–5). Although the age of haplogroup D and its geographic distribution across Eurasia roughly coincide with two important events in the cultural evolution of Eurasia—namely, the emergence and spread of domestication from the Middle East ~10,000 years ago (13) and the rapid increase in population associated with the development of cities and written language 5000 to 6000 years ago around the Middle

East (14)—the significance of this correlation is not yet clear.

References and Notes

1. J. Bond *et al.*, *Nat. Genet.* **32**, 316 (2002).
2. P. Ripoll, S. Pimpinelli, M. M. Valdivia, J. Avila, *Cell* **41**, 907 (1985).
3. J. Zhang, *Genetics* **165**, 2063 (2003).
4. P. D. Evans *et al.*, *Hum. Mol. Genet.* **13**, 489 (2004).
5. N. Kouprina *et al.*, *PLoS Biol.* **2**, E126 (2004).
6. S. L. Gilbert, W. B. Dobyns, B. T. Lahn, *Nat. Rev. Genet.* **6**, 581 (2005).
7. Materials and methods are available as supporting material on Science Online.
8. M. Bamshad, S. P. Wooding, *Nat. Rev. Genet.* **4**, 99 (2003).
9. N. A. Rosenberg *et al.*, *Science* **298**, 2381 (2002).
10. E. Zietkiewicz *et al.*, *J. Mol. Evol.* **47**, 146 (1998).
11. I. McDougall, F. H. Brown, J. G. Fleagle, *Nature* **433**, 733 (2005).
12. S. Wright, *Evolution and the Genetics of Populations* (Univ. of Chicago Press, Chicago, 1978).
13. L. L. Cavalli-Sforza, P. Menozzi, A. Piazza, *The History and Geography of Human Genes* (Princeton Univ. Press, Princeton, 1994).
14. S. D. Houston, *The First Writing: Script Invention as History and Process* (Cambridge Univ. Press, New York, 2004).
15. We thank the Coriell Institute for Medical Research, the Centre d'Etude du Polymorphisme Humain, and A. Froment for human DNA samples; and H. M. Cann, A. Di Rienzo, S. Dorus, E. E. Eichler, M. Kreitman, N. M. Pearson, and J. K. Pritchard for technical support and helpful discussions. Supported in part by the Searle Scholarship and Burroughs Wellcome Career Award (to B.T.L.) and by the David and Lucile Packard Career Award, Burroughs Wellcome Career Award, and NSF grant no. BCS-0196183 (to S.A.T.).

Supporting Online Material

www.sciencemag.org/cgi/content/full/309/5741/1720/DC1  
 Materials and Methods  
 Figs. S1 and S2  
 Tables S1 and S2  
 References and Notes

30 June 2005; accepted 10 August 2005  
 10.1126/science.1116815

# Neutral Ecological Theory Reveals Isolation and Rapid Speciation in a Biodiversity Hot Spot

Andrew M. Latimer,<sup>1\*</sup> John A. Silander Jr.,<sup>1</sup> Richard M. Cowling<sup>2</sup>

South Africa's Mediterranean-climate fynbos shrubland is a hot spot of species diversity, but its diversity patterns contrast strongly with other high-diversity areas, including the Amazon rain forest. With its extremely high levels of endemism and species turnover, fynbos is made up of dissimilar local communities that are species-rich but relatively poor in rare species. Using neutral ecological theory, we show that the relative species-abundance distributions in fynbos can be explained by migration rates that are two orders of magnitude lower than they are in tropical rain forests. Speciation rates, which are indexed by the "biodiversity parameter"  $\Theta$ , are estimated to be higher than they are in any previously examined plant system.

Regions with many endemic resident species are typically grouped together as biodiversity "hot spots," but this label obscures fundamental differences among high-diversity systems in species-diversity patterns and in the pro-

cesses that generate them. The fynbos shrubland of South Africa's Cape Floristic Region (CFR) presents a particularly strong contrast with other diverse systems such as the Amazon rain forest. Amazon rain forest tree com-

munities are dominated by species with large ranges, most of which are locally rare (1, 2), whereas most fynbos shrub species are narrow endemics, and many can be locally abundant (1, 3). Neutral ecological theory (4) enables the quantification of diversity in these contrasting systems in terms of the mechanisms that generate and partition it: rates of speciation and rates of migration.

The CFR contains over 8,000 plant species, of which a majority (69%) are endemic to the region (2). Mediterranean-climate regions are among the most diverse systems outside the tropics, and among these, the CFR has the most species in the smallest area (5). Molecular phylogenetic studies have estimated that speciation rates in some lineages in the

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of Connecticut, 75 North Eagleville Road, Storrs, CT 06269, USA. <sup>2</sup>Department of Botany, Terrestrial Ecology Research Unit, Post Office Box 77000, Nelson Mandela Metropolitan University, Port Elizabeth 6031, South Africa.

\*To whom correspondence should be addressed. E-mail: latimer@uconn.edu