

Experimental constraints on mate preferences in *Drosophila pseudoobscura* decrease offspring viability and fitness of mated pairs

Wyatt W. Anderson, Yong-Kyu Kim, and Patricia Adair Gowaty

PNAS 2007;104:4484-4488; originally published online Mar 6, 2007;
doi:10.1073/pnas.0611152104

This information is current as of March 2007.

Online Information & Services	High-resolution figures, a citation map, links to PubMed and Google Scholar, etc., can be found at: www.pnas.org/cgi/content/full/104/11/4484
References	This article cites 8 articles, 3 of which you can access for free at: www.pnas.org/cgi/content/full/104/11/4484#BIBL This article has been cited by other articles: www.pnas.org/cgi/content/full/104/11/4484#otherarticles
E-mail Alerts	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here .
Rights & Permissions	To reproduce this article in part (figures, tables) or in entirety, see: www.pnas.org/misc/rightperm.shtml
Reprints	To order reprints, see: www.pnas.org/misc/reprints.shtml

Notes:

Experimental constraints on mate preferences in *Drosophila pseudoobscura* decrease offspring viability and fitness of mated pairs

Wyatt W. Anderson*, Yong-Kyu Kim*, and Patricia Adair Gowaty†‡

*Department of Genetics, †Institute of Ecology, University of Georgia, Athens, GA 30602-2602

Contributed by Wyatt W. Anderson, December 21, 2006 (sent for review June 1, 2006)

Using *Drosophila pseudoobscura*, we tested the hypothesis that social constraints on the free expression of mate preferences, by both females and males, decrease offspring viability and reproductive success of mating pairs. Mate preference arenas eliminated intrasexual combat and intersexual coercion. The time female and male choosers spent in arena tests near either of two opposite-sex individuals measured the preferences of choosers. We placed choosers in breeding trials with their preferred or nonpreferred discriminatee when they met the minimum criteria for showing the same preference in two consecutive tests. There was no statistically significant difference in the frequency of female and male choosers meeting minimal preference criteria. There was a significant difference between female and male choosers for offspring viability, with female choice having the greater effect, but there was not a significant difference in the overall reproductive success of male and female choosers. There were significant differences in fitness between matings to preferred and nonpreferred partners. Female and male choosers paired with their nonpreferred discriminatees had offspring of significantly lower viability, as predicted by the constraints hypothesis. Reproductive success, our measure of overall fitness, was greater when males or females mated with the partner they preferred rather than the one they did not prefer.

constraints hypothesis

Social constraints on the free expression of mate preferences may affect offspring viability and reproductive success of choosers as suggested by commentators (1) on Partridge's (2) classic study, which was the first to associate offspring viability with mate choice. That mate preferences affect offspring viability also is an important prediction of hypotheses about the cues mediating mate preference (3), particularly those focused on mate compatibility at loci affecting immune function in offspring (4–6). Many studies have demonstrated that females prefer males with more elaborate secondary sexual characteristics (7), and other studies have shown that males with the most exaggerated secondary sexual traits are healthier than other males (8–12). However, relatively few studies (13–15) have shown benefits in offspring viability for females reproducing with males having traits preferred by most females. In contrast, recent studies on the benefits of mate preferences for choosers, without reference to traits of discriminatees that may mediate preferences, showed that mate preferences of females and males affect offspring viability (16–21), productivity of mating pairs (22), and chooser longevity (23). We tested the constraints hypothesis (24–26) that says social or ecological constraints on reproductive decisions, those that prevent individuals from reproducing with their preferred partner(s), result in lower offspring viability and lower overall reproductive success.

To test predictions of the constraints hypothesis, we used the same methodology with *Drosophila pseudoobscura* as Drickamer *et al.* (17, 18), Moore *et al.* (22, 23), and Bluhm and Gowaty (16, 27) did with other animals, to evaluate individual choosers' mate preferences for two opposite-sex "discriminatees" while experimentally eliminating social constraints such as intrasexual fight-

ing and intersexual coercion (Fig. 1). We picked the discriminatees at random with respect to their phenotypes to control for the possibility that signals also manipulate the decisions of choosers. That is, the experiment evaluated the effects on chooser fitness of enforced, exclusive pairing with discriminatees they did and did not prefer. Thus, this study shows the effects on the fitness of chooser individuals of an experimental constraint on the expression of mate preferences.

Results

The results of a two-way, fixed-effects ANOVA on fecundity, offspring viability, and chooser reproductive success are given in Table 1. The factors were male or female choice and also mating type, which indicated whether the chooser was paired experimentally with its preferred or nonpreferred discriminatee. Graphs of the time courses of female longevity, fecundity, offspring viability, and reproductive success by chooser sex and mating type are given in Fig. 2. There were no significant differences in female longevity between female or male chooser tests, nor were there significant differences in female longevity between matings to preferred and nonpreferred partners (Fig. 2). There were no significant differences in mean fecundity by chooser sex, mating type, or their interaction (Table 1 and Fig. 2). However, the day-by-day distribution of difference scores (preferred – nonpreferred) showed that in female choice tests, nonpreferred matings had significantly higher fecundity than did preferred matings (Fig. 3). In contrast, for male choice tests, the day-by-day distribution of difference scores (preferred – nonpreferred) for fecundity was higher in preferred matings than in nonpreferred matings (Fig. 3). Mean offspring viability did not differ significantly between male and female choice tests, and likewise, there were no significant differences in reproductive success by chooser sex (Table 1 and Fig. 2). There were statistically significant differences between pairings with preferred and nonpreferred partners for both offspring viability (Table 1 and Figs. 2 and 4) and net reproductive success of mating pairs (Table 1 and Fig. 2), both fitness measures being larger for choosers with their preferred partners. There was no indication of a significant interaction between chooser sex and mating type in either offspring viability or net reproductive success of mating pairs. In female choice tests, the net reproductive success for nonpreferred pairings was 238.08 ± 12.53 SE, and for preferred pairings, it was 286.66 ± 12.53 SE, a highly significant difference between mating types of 48.58 ± 17.72 offspring ($P < 0.01$). In male choice tests, reproductive success in nonpreferred pairings was 247.79 ± 12.04 SE, and in preferred pairings, it was 289.66 ± 12.33 SE ($P < 0.01$), a highly significant difference of 41.87 ± 17.23 SE offspring.

Author contributions: W.W.A. and P.A.G. designed research; Y.-K.K. performed research; W.W.A. and P.A.G. analyzed data; and W.W.A. and P.A.G. wrote the paper.

The authors declare no conflict of interest.

†To whom correspondence should be addressed. E-mail: gowaty@uga.edu.

© 2007 by The National Academy of Sciences of the USA




Fig. 2. Time course for female longevity, offspring viability, fecundity, and reproductive success (RS) of mating pairs in female and male choice tests, for choosers paired with their nonpreferred (filled circles) or preferred partners (open squares). Differences in longevity for females in preferred or nonpreferred matings were not statistically significant by using Kaplan–Meier statistics (for female choice tests: Wilcoxon $\chi^2 = 2.02$, $df = 1$, $P > 0.15$; for male choice tests: Wilcoxon $\chi^2 = 0.1140$, $df = 1$, $P > 0.73$).

A few drops of a live solution of yeast were added to each bottle or food cup. Twenty cups with this food medium were placed in the population cage, and on Mondays, Wednesdays, and Fridays, an old cup was replaced with a new one. Approximately 12 generations after setting up the population cage, we began separating and storing newly eclosed males and females for behavioral preference tests. We completed two large experiments in fall 1998 and fall 1999, both carried out at 22°C. We chose to do both experiments in the fall, when climatic conditions in the laboratory would be similar.

Preference Tests. We collected virgin flies from culture bottles every 7 or 8 h. We anesthetized them with CO₂, sexed them, and held them until preference testing. Flies were stored in same-sex groups of no more than 20 individuals for 7 days in vials containing food medium.

We tested the preferences of focal individuals using mate preference arenas (Fig. 1) that eliminated direct competitive interactions between discriminatees and potentially coercive

interactions between the sexes. We also used criteria for exhibiting a preference that depended on the time a chooser spent near one or the other discriminatee (see below), a methodology similar to those used by other workers for a variety of animals (7).

The design of the mate preference arenas (Fig. 1) allowed visual, olfactory, and auditory contact but prevented tactile contact between the focal individual chooser and the opposite-sex discriminatees. We picked focal individual choosers and the opposite-sex discriminatees at random with respect to any phenotypic variation obvious to us. In the female preference tests, we used 265 focal virgin females and 530 virgin male discriminatees. In the male preference tests, we used 233 focal virgin males and 466 virgin female discriminatees. We aspirated discriminatees into cells a and b, followed by aspiration of the focal chooser into corridor c.

We scored preference for one versus the other discriminatee in terms of (i) the time choosers spent in front of cells a and b and (ii) the repeatability of preference in consecutive tests using the same chooser and discriminatees. Our minimum criteria for declaring a




Fig. 3. The difference scores for fecundity between preferred and nonpreferred matings in female (A) and male (B) choice studies. The distribution of difference scores was significantly different in both female choice tests (Wilcoxon signed rank $t = -81.5$, $df = 24$, $P > |t| = 0.013$) and male choice tests (Wilcoxon signed rank $t = 59.5$, $df = 24$, $P = |t| = 0.055$). NP, nonpreferred; P, preferred.

preference in each arena test were (i) the chooser had to spend 60% (12 min) of the first 20-min preference test next to one or the other discriminatee in region a and/or b of the arena, and (ii) the chooser had to spend 60% (7.2 min) of that time with the same discriminatee (in either a or b). We switched the discriminatees between cells in consecutive tests to control for position effects. We performed female and male chooser preference tests on the same days to

control for microhabitat and temporal variation that could affect measurements of fitness components. Forty-seven percent (125 of 265) of focal female choosers and 55.3% (129 of 233) of focal male choosers met the minimum criteria for exhibiting a preference.

Breeding Tests. We discarded all individuals from preference tests in which focal choosers failed to meet the criteria for exhibiting




Fig. 4. The difference scores for offspring viability between preferred and nonpreferred matings in female (A) and male (B) choice studies. The distribution of difference scores was significantly different both for female choice tests (Wilcoxon signed rank test = 97.5, $P > |t| = 0.006$) and for male choice tests (Wilcoxon signed rank test = 134.5, $P > |t| < 0.0001$). NP, nonpreferred; P, preferred.

a preference because the hypothesis we were testing was about constraints on reproductive decisions. That is, we excluded from further tests those individuals whose preferences were ambiguous because our experiment was about the effect of constraints. When a focal chooser's associations were ambiguous, indicating no clear choice between discriminates, we could not be sure we were experimentally producing a breeding constraint when we used them in the breeding trials. We randomly assigned the first focal chooser in each series (female choosers and male choosers) to either the preferred or nonpreferred partner. Thereafter, we alternated assignments to breeding vials. We considered choosers placed with their nonpreferred discriminatee to represent greater constraint on reproductive decisions than choosers with their preferred discriminatee. The factors in our overall design thus included two experiments, sex of the chooser (female or male choice tests) and mating type (preferred or nonpreferred).

Measurements of Reproductive Success and Fitness Components. For each of the two experiments, we measured the fitness of 30–37 replicate choosers of each sex and each mating type. We measured fitness for a total of 247 pairings. We kept all pairs in mating vials for 1 day and then discarded all males. Females laid eggs from the mating that occurred with a single male after the first day of completing a preference trial. For 25 consecutive days, we moved all females to fresh vials containing food medium. We recorded longevity (l_x) as the frequency of females surviving to each day of age. We counted all eggs laid on each day, and fecundity at age x , m_x , was measured as the number of eggs each female laid during day x . Beginning 14 days after mating, we counted the number of offspring emerging as adults

from each vial over a period of 10 days. Egg-to-offspring viability at age x , v_x , was measured as the fraction of eggs laid on day x that survived to eclosion. We used reproductive success W or net reproduction (37) as our measure of the overall fitness of mating pairs. In terms of female longevity, fecundity, and offspring viability, reproductive success is $\Sigma(l_x m_x v_x)$, where the summation is over days of age.

We tested for normality of all variables. We used parametric tests throughout because all variables were normally distributed. We used a fixed-effects ANOVA with the three factors experiment, chooser sex, and mating type to test for differences among offspring viability, chooser fecundity, and chooser reproductive success. An ANOVA of our entire data set showed no significant differences between the two major experiments on any variable, so we combined the data for them. To test for independence among days for difference scores in offspring viability between preferred and nonpreferred matings, we used time series autocorrelation analyses and Fisher's Kappa to test for deviations from the null hypothesis of "white noise." We failed to reject the null hypothesis, so we concluded that the difference scores were independent between days and tested the distributions for differences using the Wilcoxon signed rank test. We considered results statistically significant if P was <0.05 .

We thank John Avise, Cindy Bluhm, Lee C. Drickamer, Steve Hubbell, Teri Markow, Allen Moore, and Jerry Downhower for many helpful and insightful conversations and for comments on the manuscript; Randy Nelson for his encouragement; Kyungsun Kim for technical assistance; and William C. Bridges for statistical advice. This work was supported by National Science Foundation Grant IBN-9631801.

- Kingett PD, Lambert DM, Telford SR (1981) *Nature* 293:492.
- Partridge L (1980) *Nature* 283:290–291.
- Hamilton WD, Zuk M (1982) *Science* 18:384–387.
- Brown JL (1997) *Behav Ecol* 8:60–65.
- Penn DJ, Potts WK (1999) *Am Nat* 153:145–164.
- Wedekind C (1999) in *Evolution in Health and Disease*, ed Stearns SC (Oxford Univ Press, Oxford), pp 102–107.
- Andersson M (1994) *Sexual Selection* (Princeton Univ Press, Princeton).
- Kacelnik A, Norris S (1998) *Behav Brain Sci* 21:378–385.
- Loehle C (1997) *Ecol Mod* 103:231–250.
- Loyau A, Saint Jalme M, Cagniant C, Sorci G (2005) *Behav Ecol Sociobiol* 58:552–557.
- Worden BD, Parker PG (2005) *Anim Behav* 70:1047–1053.
- Zala SM, Potts WK, Penn DJ (2004) *Behav Ecol* 15:338–344.
- Petrie M (1994) *Nature* 371:598–599.
- Reynolds JD, Gross MR (1992) *Proc R Soc London Ser B* 250:57–62.
- Simmons L (1987) *Behav Ecol Sociobiol* 21:313–322.
- Bluhm CK, Gowaty PA (2004) *Anim Behav* 68:977–983.
- Drickamer LC, Gowaty PA, Holmes CM (2000) *Anim Behav* 59:371–378.
- Drickamer LC, Gowaty PA, Wagner DM (2003) *Anim Behav* 65:105–114.
- Gowaty PA, Drickamer LC, Holmes S-S (2003) *Anim Behav* 65:95–103.
- Ryan KK, Altmann J (2001) *Behav Ecol Sociobiol* 50:436–440.
- Sandvik M, Rosenqvist G, Berglund A (2000) *Proc R Soc London Ser B* 267:2151–2155.
- Moore AJ, Gowaty PA, Wallin WJ, Moore PJ (2001) *Proc R Soc London Ser B* 517–523.
- Moore AJ, Gowaty PA, Moore PJ (2003) *J Evol Biol* 16:523–530.
- Gowaty PA (1996) in *Black Partnerships in Birds*, ed Black JL (Oxford Univ Press, Oxford), pp 21–52.
- Gowaty PA (2003) in *Women, Evolution, and Rape*, ed Travis C (MIT Press, Cambridge, MA), pp 61–86.
- Gowaty PA, Buschhaus N (1998) *Am Zool* 38:207–225.
- Bluhm CK, Gowaty PA (2004) *Anim Behav* 68:985–992.
- Kim Y-K, Gowaty PA, Anderson WW (2005) in *Proceedings of the Fifth International Conference on Methods and Techniques in Behavioral Research*, eds Noldus L, Grieco JFF, Loijens LWS, Zimmerman PH (Wageningen, The Netherlands), pp 533–535.
- Darwin C (1871) *The Descent of Man and Selection in Relation to Sex* (John Murray, London).
- Bateman AJ (1948) *Heredity* 2:349–368.
- Williams GC (1966) *Adaptation and Natural Selection* (Princeton Univ Press, Princeton).
- Parker GA, Baker RR, Smith VGF (1972) *J Theor Biol* 36:529–553.
- Trivers RL (1972) in *Sexual Selection and the Descent of Man*, ed Campbell B (Aldine, Chicago), pp 136–179.
- Parker GA, Simmons LW (1996) *Proc R Soc London Ser B* 315–321.
- Snook RR, Markow TA (2001) *J Insect Phys* 47:957–964.
- Snook RR, Markow TA, Karr TL (1994) *Proc Natl Acad Sci USA* 91:11222–11226.
- Futuyma DW (2005) *Evolution* (Sinauer, Sunderland, MA).